

GRAPH LIMITS AND EXCHANGEABLE RANDOM GRAPHS

PERSI DIACONIS AND SVANTE JANSON

ABSTRACT. We develop a clear connection between deFinetti's theorem for exchangeable arrays (work of Aldous–Hoover–Kallenberg) and the emerging area of graph limits (work of Lovász and many coauthors). Along the way, we translate the graph theory into more classical probability.

1. INTRODUCTION

S:intro

DeFinetti's profound contributions are now woven into many parts of probability, statistics and philosophy. Here we show how developments from deFinetti's work on partial exchangeability have a direct link to the recent development of a limiting theory for large graphs. This introduction first recalls the theory of exchangeable arrays (Section 1.1). Then, the subject of graph limits is outlined (Section 1.2). Finally, the link between these ideas, which forms the bulk of this paper, is outlined (Section 1.3).

secA

1.1. Exchangeability, partial exchangeability and exchangeable arrays. Let $\{X_i\}$, $1 \leq i < \infty$, be a sequence of binary random variables. They are *exchangeable* if

$$\mathbb{P}(X_1 = e_1, \dots, X_n = e_n) = \mathbb{P}(X_1 = e_{\sigma(1)}, \dots, X_n = e_{\sigma(n)})$$

for all n , permutations $\sigma \in \mathfrak{S}_n$ and all $e_i \in \{0, 1\}$. The celebrated representation theorem says

deF

Theorem 1.1 (deFinetti). *If $\{X_i\}$, $1 \leq i < \infty$, is a binary exchangeable sequence, then:*

- (i) *With probability 1, $X_\infty = \lim \frac{1}{n}(X_1 + \dots + X_n)$ exists.*
- (ii) *If $\mu(A) = P\{X_\infty \in A\}$, then for all n and e_i , $1 \leq i \leq n$,*

$$\mathbb{P}(X_1 = e_1, \dots, X_n = e_n) = \int_0^1 x^s (1-x)^{n-s} \mu(dx) \quad (1.1) \quad \text{eqP1}$$

for $s = e_1 + \dots + e_n$.

It is natural to refine and extend deFinetti's theorem to allow more general observables (X_i with values in a Polish space) and other notions of symmetry (partial exchangeability). A definitive treatment of these developments is

Date: December 10, 2007 (typeset January 9, 2008 16.40).

given in Kallenberg [15]. Of interest here is the extension of deFinetti's theorem to two-dimensional arrays.

Definition. Let $\{X_{ij}\}$, $1 \leq i, j < \infty$, be binary random variables. They are *separately exchangeable* if

$$\mathbb{P}(X_{ij} = e_{ij}, 1 \leq i, j \leq n) = \mathbb{P}(X_{ij} = e_{\sigma(i)\tau(j)}, 1 \leq i, j \leq n) \quad (1.2) \quad \boxed{\text{exch}}$$

for all n , all permutations $\sigma, \tau \in \mathfrak{S}_n$ and all $e_{ij} \in \{0, 1\}$. They are (*jointly*) *exchangeable* if (1.2) holds in the special case $\tau = \sigma$.

Equivalently, the array $\{X_{ij}\}$ is jointly exchangeable if the array $\{X_{\sigma(i)\sigma(j)}\}$ has the same distribution as $\{X_{ij}\}$ for every permutation σ of \mathbb{N} , and similarly for separate exchangeability.

The question of two-dimensional versions of deFinetti's theorem under (separate) exchangeability arose from the statistical problems of two-way analysis of variance. Early workers expected a version of (1.1) with perhaps a two-dimensional integral. The probabilist David Aldous [1] and the logician Douglas Hoover [14] found that the answer is more complicated.

Define a random binary array $\{X_{ij}\}$ as follows: Let U_i, V_j , $1 \leq i, j < \infty$, be independent and uniform in $[0, 1]$. Let $W(x, y)$ be a function from $[0, 1]^2$ to $[0, 1]$. Let X_{ij} be 1 or 0 as a $W(U_i, V_j)$ -coin comes up heads or tails. Let P_W be the probability distribution of $\{X_{ij}\}$, $1 \leq i, j < \infty$. The family $\{X_{ij}\}$ is separately exchangeable because of the symmetry of the construction. The Aldous–Hoover theorem says that any separately exchangeable binary array is a mixture of such P_W :

Theorem 1.2 (Aldous–Hoover). *Let $X = \{X_{ij}\}$, $1 \leq i, j < \infty$, be a separately exchangeable binary array. Then, there is a probability μ such that*

$$\mathbb{P}\{X \in A\} = \int P_W(A) \mu(dW).$$

There is a similar result for jointly exchangeable arrays.

The uniqueness of μ resisted understanding; if \widehat{W} is obtained from W by a measure-preserving change of each variable, clearly the associated process $\{\widehat{X}_{ij}\}$ has the same joint distribution as $\{X_{ij}\}$. Using model theory, Hoover [14] was able to show that this was the only source of non-uniqueness. A ‘probabilist’s proof’ was finally found by Kallenberg, see [15, Sect. 7.6] for details and references.

These results hold for higher dimensional arrays with X_{ij} taking values in a Polish space with minor change [15, Chap. 7]. The description above has not mentioned several elegant results of the theory. In particular, Kallenberg’s ‘spreadable’ version of the theory replaces invariance under a group by invariance under subsequences. A variety of tail fields may be introduced to allow characterizing when W takes values in $\{0, 1\}$ [10, Sect. 4]. Much more general notions of partial exchangeability are studied in [11].

secB

1.2. Graph limits. Large graphs, both random and deterministic, abound in applications. They arise from the internet, social networks, gene regulation, ecology and in mathematics. It is natural to seek an approximation theory: What does it mean for a sequence of graphs to converge? When can a large complex graph be approximated by a small graph?

In a sequence of papers [6, 7, 8, 9, 13, 16, 17, 18, 21, 20, 22, 19], Laszlo Lovász with coauthors (listed here in order of frequency) V. T. Sós, B. Szegedy, C. Borgs, J. Chayes, K. Vesztergombi, A. Schrijver, M. Freedman have developed a beautiful, unifying limit theory. This sheds light on topics such as graph homomorphisms, Szemerédi’s regularity lemma, quasi-random graphs, graph testing and extremal graph theory. Their theory has been developed for dense graphs (number of edges comparable with the square of number of vertices) but parallel theories for sparse graphs are beginning to emerge [4].

Roughly, a growing sequence of finite graphs G_n converges if, for any fixed graph F , the proportion of copies of F in G_n converges. Section 2 below has precise definitions.

Example 1.3. Define a probability distribution on graphs on n -vertices as follows. Flip a θ -coin for each vertex (dividing vertices into ‘boys’ and ‘girls’). Connect two boys with probability p . Connect two girls with probability p' . Connect a boy and a girl with probability p'' . Thus, if $p = p' = 0, p'' = 1$, we have a random bipartite graph. If $p = p' = 1, p'' = 0$, we have two disjoint complete graphs. If $p = p' = p''$, we have the Erdős–Renyi model. As n grows, these models generate a sequence of random graphs which converge almost surely to a limiting object described below.

More substantial examples involving random threshold graphs are in [12]. SJ

If a sequence of graphs converges, what does it converge to? For exchangeable random graphs (defined below), there is a limiting object which may be thought of as a probability measure on infinite random graphs. Suppose $W(x, y) = W(y, x)$ is a function from $[0, 1]^2 \rightarrow [0, 1]$. Choose $\{U_i\}$, $1 \leq i < \infty$, independent uniformly distributed random variables on $[0, 1]$. Form an infinite random graph by putting an edge from i to j with probability $W(U_i, U_j)$. This measure on graphs (or alternatively W) is the limiting object.

For the “boys and girls” example above, W may be pictured as

	θ	p	p''	
		p''	p'	
	0	θ	1	

The theory developed shows that various properties of G_n can be well approximated by calculations with the limiting object. There is an elegant characterization of these ‘continuous graph properties’ with applications to algorithms for graph testing (Does this graph contain an Eulerian cycle?) or parameter estimation (What is an approximation to the size of the maximum cut?). There is a practical way to find useful approximations to a large graph by graphs of fixed size [6]. This paper also contains a useful review of the current state of the theory with proofs and references.

We have sketched the theory for unweighted graphs. There are generalizations to graphs with weights on vertices and edges, to bipartite, directed and hypergraphs. The sketch leaves out many nice developments. For example, the useful cut metric between graphs [19] and connections to statistical physics [9].

secC

1.3. Overview of the present paper. There is an apparent similarity between the measure P_W of the Aldous–Hoover theorem and the limiting object W from graph limits. Roughly, working with symmetric W gives the graph limit theory; working with general W gives directed graphs. The main results of this paper make these connections precise.

Basic definitions are in Section 2 which introduces a probabilist’s version of graph convergence equivalent to the definition using graph homomorphisms. Section 3 uses the well-established theory of weak convergence of a sequence of probability measures on a metric space to get properties of graph convergence. Section 4 carries things over to infinite graphs.

The main results appear in Section 5. This introduces exchangeable random graphs and gives a one-to-one correspondence between infinite exchangeable random graphs and distributions on the space of proper graph limits (Theorem 5.3), which specializes to a one-to-one correspondence between proper graph limits and extreme points in the set of distributions of exchangeable random graphs (Corollary 5.4).

A useful characterization of the extreme points of the set of exchangeable random graphs is in Theorem 5.5. These results are translated to the equivalence between proper graph limits and the Aldous–Hoover theory in Section 6. The non-uniqueness of the representing W , for exchangeable random graphs and for graph limits, is discussed in Section 7.

The equivalence involves symmetric $W(x, y)$ and a single permutation σ taking $W(U_i, U_j)$ to $W(U_{\sigma(i)}, U_{\sigma(j)})$. The original Aldous–Hoover theorem, with perhaps non-symmetric $W(x, y)$ and $W(U_i, V_j)$ to $W(U_{\sigma(i)}, V_{\tau(j)})$ translates to a limit theorem for bipartite graphs. This is developed in Section 8. The third case of the Aldous–Hoover theory for two-dimensional arrays, perhaps non-symmetric $W(x, y)$ and a single permutation σ , corresponds to directed graphs; this is sketched in Section 9.

The extensions to weighted graphs are covered by allowing X_{ij} to take general values in the Aldous–Hoover theory. The extension to hypergraphs

SJ

SJ

SJ

follows from the Aldous–Hoover theory for higher-dimensional arrays. (The details of these extensions are left to the reader.)

SJ

Despite these parallels, the theories have much to contribute to each other. The algorithmic, graph testing, Szemerédi partitioning perspective is new to exchangeability theory. Indeed, the “boys and girls” random graph was introduced to study the psychology of vision in Diaconis–Freedman (1981). As far as we know, its graph theoretic properties have not been studied. The various developments around shell-fields in exchangeability, which characterize zero/one $W(x, y)$, have yet to be translated into graph-theoretic terms.

Acknowledgements. This lecture is an extended version of a talk presented by PD at the 100th anniversary of deFinetti’s birth in Rome, 2006. We thank the organizers.

SJ thanks Christian Borgs and Jennifer Chayes for inspiration from lectures and discussions during the Oberwolfach meeting ‘Combinatorics, Probability and Computing’, held in November, 2006. Parts of the research were completed during a visit by SJ to the Université de Nice - Sophia Antipolis in January 2007.

2. DEFINITIONS AND BASIC PROPERTIES

Sdef

All graphs will be simple. Infinite graphs will be important in later sections, but will always be clearly stated to be infinite; otherwise, graphs will be finite. We denote the vertex and edge sets of a graph G by $V(G)$ and $E(G)$, and the numbers of vertices and edges by $v(G) := |V(G)|$ and $e(G) := |E(G)|$. We consider both labelled and unlabelled graphs; the labels will be the integers $1, \dots, n$, where n is the number of vertices in the graph. A labelled graph is thus a graph with vertex set $[n] := \{1, \dots, n\}$ for some $n \geq 1$; we let \mathcal{L}_n denote the set of the $2^{\binom{n}{2}}$ labelled graphs on $[n]$ and let $\mathcal{L} := \bigcup_{n=1}^{\infty} \mathcal{L}_n$. An unlabelled graph can be regarded as a labelled graph where we ignore the labels; formally, we define \mathcal{U}_n , the set of unlabelled graphs of order n , as the quotient set \mathcal{L}_n / \cong of labelled graphs modulo isomorphisms. We let $\mathcal{U} := \bigcup_{n=1}^{\infty} \mathcal{U}_n = \mathcal{L} / \cong$, the set of all unlabelled graphs.

Note that we can, and often will, regard a labelled graph as an unlabelled graph.

If G is an (unlabelled) graph and v_1, \dots, v_k is a sequence of vertices in G , then $G(v_1, \dots, v_k)$ denotes the labelled graph with vertex set $[k]$ where we put an edge between i and j if v_i and v_j are adjacent in G . We allow the possibility that $v_i = v_j$ for some i and j . (In this case, there is no edge ij because there are no loops in G .)

We let $G[k]$, for $k \geq 1$, be the random graph $G(v_1, \dots, v_k)$ obtained by sampling v_1, \dots, v_k uniformly at random among the vertices of G , with replacement. In other words, v_1, \dots, v_k are independent uniformly distributed random vertices of G .

For $k \leq v(G)$, we further let $G[k]'$ be the random graph $G(v'_1, \dots, v'_k)$ where we sample v'_1, \dots, v'_k uniformly at random without replacement; the

sequence v'_1, \dots, v'_k is thus a uniformly distributed random sequence of k distinct vertices.

The graph limit theory in [19] and subsequent papers is based on the study of the functional $t(F, G)$ which is defined for two graphs F and G as the proportion of all mappings $V(F) \rightarrow V(G)$ that are graph homomorphisms $F \rightarrow G$, i.e., map adjacent vertices to adjacent vertices. In probabilistic terms, $t(F, G)$ is the probability that a uniform random mapping $V(F) \rightarrow V(G)$ is a graph homomorphism. Using the notation introduced above, we can, equivalently, write this as, assuming that F is labelled and $k = v(F)$,

$$t(F, G) := \mathbb{P}(F \subseteq G[k]). \quad (2.1) \quad \boxed{\text{t}}$$

Note that both F and $G[k]$ are graphs on $[k]$, so the relation $F \subseteq G[k]$ is well-defined as containment of labelled graphs on the same vertex set, i.e. as $E(F) \subseteq E(G[k])$. Although the relation $F \subseteq G[k]$ may depend on the labelling of F , the probability in (2.1) does not, by symmetry, so $t(F, G)$ is really well defined by (2.1) for unlabelled F and G .

With F, G and k as in (2.1), we further define, again following [19] (and the notation of [8]) but stating the definitions in different but equivalent forms,

$$t_{\text{inj}}(F, G) := \mathbb{P}(F \subseteq G[k]') \quad (2.2) \quad \boxed{\text{t}_{\text{inj}}}$$

and

$$t_{\text{ind}}(F, G) := \mathbb{P}(F = G[k]'), \quad (2.3) \quad \boxed{\text{t}_{\text{ind}}}$$

provided F and G are (unlabelled) graphs with $v(F) \leq v(G)$. If $v(F) > v(G)$ we set $t_{\text{inj}}(F, G) := t_{\text{ind}}(F, G) := 0$.

Since the probability that a random sample v_1, \dots, v_k of vertices in G contains some repeated vertex is $\leq k^2/(2v(G))$, it follows that [19]

$$|t(F, G) - t_{\text{inj}}(F, G)| \leq \frac{v(F)^2}{2v(G)}. \quad (2.4) \quad \boxed{\text{a4}}$$

Hence, when considering asymptotics with $v(G) \rightarrow \infty$, it does not matter whether we use t or t_{inj} . Moreover, if $F \in \mathcal{L}_k$, then, as pointed out in [8] and [19],

$$t_{\text{inj}}(F, G) = \sum_{F' \in \mathcal{L}_k, F' \supseteq F} t_{\text{ind}}(F, G) \quad (2.5) \quad \boxed{\text{a4a}}$$

and, by inclusion-exclusion,

$$t_{\text{ind}}(F, G) = \sum_{F' \in \mathcal{L}_k, F' \supseteq F} (-1)^{e(F') - e(F)} t_{\text{inj}}(F, G). \quad (2.6) \quad \boxed{\text{a4b}}$$

Hence, the two families $\{t_{\text{inj}}(F, \cdot)\}_{F \in \mathcal{U}}$ and $\{t_{\text{ind}}(F, \cdot)\}_{F \in \mathcal{U}}$ of graph functionals contain the same information and can replace each other.

The basic definition of Lovász and Szegedy [19] and Borgs, Chayes, Lovász, Sós and Vesztergombi [8] is that a sequence (G_n) of graphs converges if

$t(F, G_n)$ converges for every graph F . We can express this by considering the map $\tau : \mathcal{U} \rightarrow [0, 1]^{\mathcal{U}}$ defined by

$$\tau(G) := (t(F, G))_{F \in \mathcal{U}} \in [0, 1]^{\mathcal{U}}. \tag{2.7} \quad \boxed{\text{tau}}$$

Then (G_n) converges if and only if $\tau(G_n)$ converges in $[0, 1]^{\mathcal{U}}$, equipped with the usual product topology. Note that $[0, 1]^{\mathcal{U}}$ is a compact metric space; as is well known, a metric can be defined by, for example,

$$d((x_F), (y_F)) := \sum_{i=0}^{\infty} 2^{-i} |x_{F_i} - y_{F_i}|, \tag{2.8} \quad \boxed{\text{a5}}$$

where F_1, F_2, \dots is some enumeration of all unlabelled graphs.

We define $\mathcal{U}^* := \tau(\mathcal{U}) \subseteq [0, 1]^{\mathcal{U}}$ to be the image of \mathcal{U} under this mapping τ , and let $\overline{\mathcal{U}^*}$ be the closure of \mathcal{U}^* in $[0, 1]^{\mathcal{U}}$. Thus $\overline{\mathcal{U}^*}$ is a compact metric space. (For explicit descriptions of the subset $\overline{\mathcal{U}^*}$ of $[0, 1]^{\mathcal{U}}$ as a set of graph functionals, see Lovász and Szegedy [19].)

As pointed out in [19] and [8] (in equivalent terminology), τ is not injective; for example, $\tau(K_{n,n})$ is the same for all complete bipartite graphs $K_{n,n}$. Nevertheless, as in [19] and [8], we can consider a graph G as an element of \mathcal{U}^* by identifying G and $\tau(G)$ (thus identifying graphs with the same $\tau(G)$), and then convergence of (G_n) as defined above is equivalent to convergence in $\overline{\mathcal{U}^*}$. The limit is thus an element of $\overline{\mathcal{U}^*}$, but typically not a graph in \mathcal{U}^* . The main result of Lovász and Szegedy [19] is a representation of the elements in $\overline{\mathcal{U}^*}$ to which we will return in Section 6.

Rmetric1

Remark 2.1. As said above, $\overline{\mathcal{U}^*}$ is a compact metric space, and it can be given several equivalent metrics. One metric is the metric (2.8) inherited from $[0, 1]^{\mathcal{U}}$, which for graphs becomes $d(G, G') = \sum_i 2^{-i} |t(F_i, G) - t(F_i, G')|$. Another metric, shown by Borgs, Chayes, Lovász, Sós and Vesztergombi [8] to be equivalent, is the cut-distance δ_{\square} , see [8] for definitions. Further characterizations of convergence of sequences of graphs in $\overline{\mathcal{U}}$ are given in Borgs, Chayes, Lovász, Sós and Vesztergombi [8, 9].

SJ

The identification of graphs with the same image in \mathcal{U}^* (i.e., with the same $t(F, \cdot)$ for all F) is sometimes elegant but at other times inconvenient. It can be avoided if we instead let \mathcal{U}^+ be the union of \mathcal{U} and some one-point set $\{*\}$ and consider the mapping $\tau^+ : \mathcal{U} \rightarrow [0, 1]^{\mathcal{U}^+} = [0, 1]^{\mathcal{U}} \times [0, 1]$ defined by

$$\tau^+(G) = (\tau(G), v(G)^{-1}). \tag{2.9} \quad \boxed{\text{tau+}}$$

Then τ^+ is injective, because if $\tau(G_1) = \tau(G_2)$ for two graphs G_1 and G_2 with the same number of vertices, then G_1 and G_2 are isomorphic and thus $G_1 = G_2$ as unlabelled graphs. (This can easily be shown directly: it follows from (2.1) that $G_1[k] \stackrel{d}{=} G_2[k]$ for every k , which implies $G_1[k]' \stackrel{d}{=} G_2[k]'$ for every $k \leq v(G_1) = v(G_2)$; now take $k = v(G_1)$. It is also a consequence of [8, Theorem 2.7 and Theorem 2.3 or Lemma 5.1].)

Consequently, we can identify \mathcal{U} with its image $\tau^+(\mathcal{U}) \subseteq [0, 1]^{\mathcal{U}^+}$ and define $\overline{\mathcal{U}} \subseteq [0, 1]^{\mathcal{U}^+}$ as its closure. It is easily seen that a sequence (G_n) of graphs converges in $\overline{\mathcal{U}}$ if and only if either $v(G_n) \rightarrow \infty$ and (G_n) converges in $\overline{\mathcal{U}^*}$, or the sequence (G_n) is constant from some n_0 on. Hence, convergence in $\overline{\mathcal{U}}$ is essentially the same as the convergence considered by Lovász and Szegedy [19], but without any identification of non-isomorphic graphs of different orders.

Alternatively, we can consider τ_{inj} or τ_{ind} defined by

$$\begin{aligned}\tau_{\text{inj}}(G) &:= (t_{\text{inj}}(F, G))_{F \in \mathcal{U}} \in [0, 1]^{\mathcal{U}}, \\ \tau_{\text{ind}}(G) &:= (t_{\text{ind}}(F, G))_{F \in \mathcal{U}} \in [0, 1]^{\mathcal{U}}.\end{aligned}$$

It is easy to see that both τ_{inj} and τ_{ind} are injective mappings $\mathcal{U} \rightarrow [0, 1]^{\mathcal{U}}$. (If $\tau_{\text{inj}}(F, G_1) = \tau_{\text{inj}}(F, G_2)$ for all F , we take $F = G_1$ and $F = G_2$ and conclude $G_1 = G_2$, using our special definition above when $v(F) > v(G)$.) Hence, we can again identify \mathcal{U} with its image and consider its closure $\overline{\mathcal{U}}$ in $[0, 1]^{\mathcal{U}}$. Moreover, using (2.4), (2.5), and (2.6), it is easily shown that if (G_n) is a sequence of unlabelled graphs, then

$$\tau^+(G_n) \text{ converges} \iff \tau_{\text{ind}}(G_n) \text{ converges} \iff \tau_{\text{inj}}(G_n) \text{ converges}.$$

Hence, the three compactifications $\overline{\tau^+(\mathcal{U})}$, $\overline{\tau_{\text{inj}}(\mathcal{U})}$, $\overline{\tau_{\text{ind}}(\mathcal{U})}$ are homeomorphic and we can use any of them for $\overline{\mathcal{U}}$. We let $\mathcal{U}_\infty := \overline{\mathcal{U}} \setminus \mathcal{U}$; this is the set of all limit objects of sequences (G_n) in \mathcal{U} with $v(G_n) \rightarrow \infty$. (I.e., it is the set of all proper graph limits.)

We will in the sequel prefer to use $\overline{\mathcal{U}}$ rather than $\overline{\mathcal{U}^*}$, thus not identifying some graphs of different orders, nor identifying finite graphs with some limit objects in \mathcal{U}_∞ .

For every fixed graph F , the functions $t(F, \cdot)$, $t_{\text{inj}}(F, \cdot)$ and $t_{\text{ind}}(F, \cdot)$ have unique continuous extensions to $\overline{\mathcal{U}}$, for which we use the same notation. We similarly extend $v(\cdot)^{-1}$ continuously to $\overline{\mathcal{U}}$ by defining $v(G) = \infty$ and thus $v(G)^{-1} = 0$ for $G \in \mathcal{U}_\infty := \overline{\mathcal{U}} \setminus \mathcal{U}$. Then (2.4), (2.5) and (2.6) hold for all $G \in \overline{\mathcal{U}}$, where (2.4) means that

$$t_{\text{inj}}(F, G) = t(F, G), \quad G \in \mathcal{U}_\infty. \quad (2.10) \quad \boxed{\text{a4x}}$$

Note that $\overline{\mathcal{U}}$ is a compact metric space. Different, equivalent, metrics are given by the embeddings τ^+ , τ_{inj} , τ_{ind} into $[0, 1]^{\mathcal{U}^+}$ and $[0, 1]^{\mathcal{U}}$. Another equivalent metric is, by Remark 2.1 and the definition of τ^+ , $\delta_\square(G_1, G_2) + |v(G_1)^{-1} - v(G_2)^{-1}|$.

We summarize the results above on convergence.

T1 **Theorem 2.1.** *A sequence (G_n) of graphs converges in the sense of Lovász and Szegedy [19] if and only if it converges in the compact metric space $\overline{\mathcal{U}^*}$. Moreover, if $v(G_n) \rightarrow \infty$, the sequence (G_n) converges in this sense if and only if it converges in $\overline{\mathcal{U}}$.*

The projection $\pi : [0, 1]^{\mathcal{U}^+} = [0, 1]^{\mathcal{U}} \times [0, 1] \rightarrow [0, 1]^{\mathcal{U}}$ maps $\tau^+(G)$ to $\tau(G)$ for every graph G , so by continuity it maps $\overline{\mathcal{U}}$ into $\overline{\mathcal{U}^*}$. For graph $G \in \mathcal{U}$, $\pi(G) = \tau(G)$ is the object in $\overline{\mathcal{U}^*}$ corresponding to G considered above, and we will in the sequel denote this object by $\pi(G)$; recall that this projection $\mathcal{U} \rightarrow \overline{\mathcal{U}^*}$ is not injective. (We thus distinguish between a graph G and its “ghost” $\pi(G)$ in $\overline{\mathcal{U}^*}$. Recall that when graphs are considered as elements of $\overline{\mathcal{U}^*}$ as in [19] and [8], certain graphs are identified with each other; we avoid this.) On the other hand, an element G of $\overline{\mathcal{U}}$ is by definition determined by $\tau(G)$ and $v(G)^{-1}$, cf. (2.9), so the restriction $\pi : \mathcal{U}_n \rightarrow \overline{\mathcal{U}^*}$ is injective for each $n \leq \infty$. In particular, $\pi : \mathcal{U}_\infty \rightarrow \overline{\mathcal{U}^*}$ is injective. Moreover, this map is surjective because every element $G \in \overline{\mathcal{U}^*}$ is the limit of some sequence (G_n) of graphs in \mathcal{U} with $v(G_n) \rightarrow \infty$; by Theorem 2.1, this sequence converges in $\overline{\mathcal{U}}$ to some element G' , and then $\pi(G') = G$. Since \mathcal{U}_∞ is compact, the restriction of π to \mathcal{U}_∞ is thus a homeomorphism, and we have the following theorem, saying that we can identify the set \mathcal{U}_∞ of proper graph limits with $\overline{\mathcal{U}^*}$.

Tcuoo **Theorem 2.2.** *The projection π maps the set $\mathcal{U}_\infty := \overline{\mathcal{U}} \setminus \mathcal{U}$ of proper graph limits homeomorphically onto $\overline{\mathcal{U}^*}$.*

3. CONVERGENCE OF RANDOM GRAPHS

Sconv

A *random unlabelled graph* is a random element of \mathcal{U} (with any distribution; we do not imply any particular model). We consider convergence of a sequence (G_n) of random unlabelled graphs in the larger space \mathcal{U} ; recall that this is a compact metric space so we may use the general theory set forth in, for example, Billingsley [2].

We use the standard notations \xrightarrow{d} , \xrightarrow{p} , $\xrightarrow{\text{a.s.}}$ for convergence in distribution, probability, and almost surely, respectively. We will only consider the case when $v(G_n) \rightarrow \infty$, at least in probability. (The reader may think of the case when G_n has n vertices, although that is not necessary in general.)

We begin with convergence in distribution.

T2 **Theorem 3.1.** *Let G_n , $n \geq 1$, be random unlabelled graphs and assume that $v(G_n) \xrightarrow{p} \infty$. The following are equivalent, as $n \rightarrow \infty$.*

- T2a** (i) $G_n \xrightarrow{d} \Gamma$ for some random $\Gamma \in \overline{\mathcal{U}}$.
- T2b** (ii) For every finite family F_1, \dots, F_m of (non-random) graphs, the random variables $t(F_1, G_n), \dots, t(F_m, G_n)$ converge jointly in distribution.
- T2c** (iii) For every (non-random) $F \in \mathcal{U}$, the random variables $t(F, G_n)$ converge in distribution.
- T2d** (iv) For every (non-random) $F \in \mathcal{U}$, the expectations $\mathbb{E}t(F, G_n)$ converge.

If these properties hold, then the limits in (ii), (iii) and (iv) are $(t(F_i, \Gamma))_{i=1}^m$, $t(F, \Gamma)$ and $\mathbb{E}t(F, \Gamma)$, respectively. Furthermore, $\Gamma \in \mathcal{U}_\infty$ a.s.

The same results hold if t is replaced by t_{inj} or t_{ind} .

Proof. (i) \iff (ii). Since $\overline{\mathcal{U}}$ is a closed subset of $[0, 1]^{\mathcal{U}^+}$, convergence in distribution in $\overline{\mathcal{U}}$ is equivalent to convergence of $\tau^+(G_n) = ((t(F, G_n))_{F \in \mathcal{U}}, v(G_n)^{-1})$ in $[0, 1]^{\mathcal{U}^+}$. Since we assume $v(G_n)^{-1} \xrightarrow{\text{P}} 0$, this is equivalent to convergence of $(t(F, G_n))_{F \in \mathcal{U}}$ in $[0, 1]^{\mathcal{U}}$ [2, Theorem 4.4], which is equivalent to convergence in distribution of all finite families $(t(F_i, G_n))_{i=1}^m$.

(ii) \implies (iii). Trivial.

(iii) \implies (iv). Immediate, since t is bounded (by 1).

(iv) \implies (ii). Let F_1, \dots, F_m be fixed graphs and let ℓ_1, \dots, ℓ_m be positive integers. Let F be the disjoint union of ℓ_i copies of F_i , $i = 1, \dots, m$. Then, for every $G \in \mathcal{U}$, from the definition of t ,

$$t(F, G) = \prod_{i=1}^m t(F_i, G)^{\ell_i},$$

and hence

$$\mathbb{E} \prod_{i=1}^m t(F_i, G)^{\ell_i} = \mathbb{E} t(F, G). \quad (3.1) \quad \boxed{\text{b3}}$$

Consequently, if (iv) holds, then every joint moment $\mathbb{E} \prod_{i=1}^m t(F_i, G)^{\ell_i}$ of $t(F_1, G_n), \dots, t(F_m, G_n)$ converges. Since $t(F_i, G_n)$ are bounded (by 1), this implies joint convergence in distribution by the method of moments.

The identification of the limits is immediate. Since $v(G_n) \xrightarrow{\text{P}} \infty$, (i) implies that $v(\Gamma) = \infty$ a.s., and thus $\Gamma \in \mathcal{U}_\infty$.

Finally, it follows from (2.4), (2.5) and (2.6) that we can replace t by t_{inj} or t_{ind} in (ii) and (iv), and the implications (ii) \implies (iii) and (iii) \implies (iv) are immediate for t_{inj} and t_{ind} too. \square

Specializing to the case of a non-random limit $G \in \mathcal{U}_\infty$, we obtain the corresponding result for convergence in probability.

$\boxed{\text{C2}}$ **Corollary 3.2.** *Let $G_n, n \geq 1$, be random unlabelled graphs such that $v(G_n) \xrightarrow{\text{P}} \infty$, and let $G \in \mathcal{U}_\infty$. The following are equivalent, as $n \rightarrow \infty$.*

- $\boxed{\text{C2a}}$ (i) $G_n \xrightarrow{\text{P}} G$.
- $\boxed{\text{C2c}}$ (ii) $t(F, G_n) \xrightarrow{\text{P}} t(F, G)$ for every (non-random) $F \in \mathcal{U}$.
- $\boxed{\text{C2d}}$ (iii) $\mathbb{E} t(F, G_n) \rightarrow t(F, G)$ for every (non-random) $F \in \mathcal{U}$.

The same result holds if t is replaced by t_{inj} or t_{ind} .

Note further that under the same assumptions, it follows directly from Theorem 2.1 that $G_n \xrightarrow{\text{a.s.}} G$ if and only if $t(F, G_n) \xrightarrow{\text{a.s.}} t(F, G)$ for every $F \in \mathcal{U}$.

We observe another corollary to Theorem 3.1 (and its proof).

C2aa

Corollary 3.3. *If Γ is a random element of $\mathcal{U}_\infty = \overline{\mathcal{U}} \setminus \mathcal{U} \cong \overline{\mathcal{U}^*}$, then, for every sequence F_1, \dots, F_m of graphs, possibly with repetitions,*

$$\mathbb{E} \prod_{i=1}^m t(F_i, \Gamma) = \mathbb{E} t(\oplus_{i=1}^m F_i, \Gamma), \tag{3.2} \quad \text{b3x}$$

where $\oplus_{i=1}^m F_i$ denotes the disjoint union of F_1, \dots, F_m . As a consequence, the distribution of Γ is uniquely determined by the numbers $\mathbb{E} t(F, \Gamma)$, $F \in \mathcal{U}$. Alternatively, the distribution of Γ is uniquely determined by the numbers $\mathbb{E} t_{\text{ind}}(F, \Gamma)$, $F \in \mathcal{U}$.

Proof. Since \mathcal{U} is dense in $\overline{\mathcal{U}} \supseteq \mathcal{U}_\infty$, there exists random unlabelled graphs G_n such that $G_n \xrightarrow{\text{a.s.}} \Gamma$. In particular, $G_n \xrightarrow{d} \Gamma$ and $v(G_n) \xrightarrow{p} \infty$ (in fact, we may assume $v(G_n) = n$), so Theorem 3.1 and its proof apply, and (3.2) follows from (3.1) applied to G_n by letting $n \rightarrow \infty$.

For the final statement, note that (3.2) shows that the expectations $\mathbb{E} t(F, \Gamma)$, $F \in \mathcal{U}$, determine all moments $\mathbb{E} \prod_{i=1}^m t(F_i, \Gamma)$, and thus the joint distribution of $t(F, \Gamma)$, $F \in \mathcal{U}$, which is the same as the distribution of $\tau(\Gamma) = (t(F, \Gamma))_{F \in \mathcal{U}} \in [0, 1]^{\mathcal{U}}$, and we have defined \mathcal{U}_∞ such that we identify Γ and $\tau(\Gamma)$. Finally, the numbers $\mathbb{E} t_{\text{ind}}(F, \Gamma)$, $F \in \mathcal{U}$, determine all $\mathbb{E} t(F, \Gamma)$ by (2.5), recalling that $t_{\text{inj}}(F, \Gamma) = t(F, \Gamma)$ by (2.10). \square

Remark 3.1. The numbers $\mathbb{E} t(F, \Gamma)$ for a random $\Gamma \in \mathcal{U}_\infty$ thus play a role similar to the one played by moments for a random variable. (And the relation between $\mathbb{E} t(F, \Gamma)$ and $\mathbb{E} t_{\text{ind}}(F, \Gamma)$ has some resemblance to the relation between moments and cumulants.)

4. CONVERGENCE TO INFINITE GRAPHS

Sinfinite

We will in this section consider also labelled *infinite* graphs with the vertex set $\mathbb{N} = \{1, 2, \dots\}$. Let \mathcal{L}_∞ denote the set of all such graphs. These graphs are determined by their edge sets, so \mathcal{L}_∞ can be identified with the power set $\mathcal{P}(E(K_\infty))$ of all subsets of the edge set $E(K_\infty)$ of the complete infinite graph K_∞ , and thus with the infinite product set $\{0, 1\}^{E(K_\infty)}$. We give this space, and thus \mathcal{L}_∞ , the product topology. Hence, \mathcal{L}_∞ is a compact metric space.

It is sometimes convenient to regard \mathcal{L}_n for a finite n as a subset of \mathcal{L}_∞ : we can identify graphs in \mathcal{L}_n and \mathcal{L}_∞ with the same edge set. In other words, if $G \in \mathcal{L}_n$ is a graph with vertex set $[n]$, we add an infinite number of isolated vertices $n + 1, n + 2, \dots$ to obtain a graph in \mathcal{L}_∞ .

Conversely, if $H \in \mathcal{L}_\infty$ is an infinite graph, we let $H|_{[n]} \in \mathcal{L}_n$ be the induced subgraph of H with vertex set $[n]$.

If G is a (finite) graph, let \widehat{G} be the random labelled graph obtained by a random labelling of the vertices of G by the numbers $1, \dots, v(G)$. (If G is labelled, we thus ignore the labels and randomly relabel.) Thus \widehat{G} is a random finite graph with the same number of vertices as G , but as just said, we can (and will) also regard \widehat{G} as a random graph in \mathcal{L}_∞ .

We use the same notation \widehat{G} also for a random (finite) graph G given a random labelling.

TC1

Theorem 4.1. *Let (G_n) be a sequence of random graphs in \mathcal{U} and assume that $v(G_n) \xrightarrow{P} \infty$. Then the following are equivalent.*

- (i) $G_n \xrightarrow{d} \Gamma$ in $\overline{\mathcal{U}}$ for some random $\Gamma \in \overline{\mathcal{U}}$.
- (ii) $\widehat{G}_n \xrightarrow{d} H$ in \mathcal{L}_∞ for some random $H \in \mathcal{L}_\infty$.

If these hold, then $\mathbb{P}(H|_{[k]} = F) = \mathbb{E} t_{\text{ind}}(F, \Gamma)$ for every $F \in \mathcal{L}_k$. Furthermore, $\Gamma \in \mathcal{U}_\infty$ a.s.

Proof. Let G be a labelled graph and consider the graph $\widehat{G}|_{[k]}$, assuming $k \leq v(G)$. This random graph equals $G[k]' = G(v'_1, \dots, v'_k)$, where v'_1, \dots, v'_k are k vertices sampled at random without replacement as in Section 2. Hence, by (2.3), for every $F \in \mathcal{L}_k$,

$$\mathbb{P}(\widehat{G}|_{[k]} = F) = t_{\text{ind}}(F, G), \quad \text{if } k \leq v(G).$$

Applied to the random graph G_n , this yields

$$\mathbb{E} t_{\text{ind}}(F, G_n) \leq \mathbb{P}(\widehat{G}_n|_{[k]} = F) \leq \mathbb{E} t_{\text{ind}}(F, G_n) + P(v(G_n) < k). \quad (4.1) \quad \text{c3}$$

By assumption, $P(v(G_n) < k) \rightarrow 0$ as $n \rightarrow \infty$, and it follows from (4.1) and Theorem 3.1 that $G_n \xrightarrow{d} \Gamma$ in $\overline{\mathcal{U}}$ if and only if

$$\mathbb{P}(\widehat{G}_n|_{[k]} = F) \rightarrow \mathbb{E} t_{\text{ind}}(F, \Gamma) \quad (4.2) \quad \text{c3a}$$

for every $k \geq 1$ and every $F \in \mathcal{L}_k$.

Since \mathcal{L}_k is a finite set, (4.2) says that, for every k , $\widehat{G}_n|_{[k]} \xrightarrow{d} H_k$ for some random graph $H_k \in \mathcal{L}_k$ with $\mathbb{P}(H_k = F) = \mathbb{E} t_{\text{ind}}(F, \Gamma)$ for $F \in \mathcal{L}_k$. Since \mathcal{L}_∞ has the product topology, this implies $\widehat{G}_n \xrightarrow{d} H$ in \mathcal{L}_∞ for some random $H \in \mathcal{L}_\infty$ with $H|_{[k]} \stackrel{d}{=} H_k$.

Conversely, if $\widehat{G}_n \xrightarrow{d} H$ in \mathcal{L}_∞ , then $\widehat{G}_n|_{[k]} \xrightarrow{d} H|_{[k]}$ so the argument above shows that

$$\mathbb{E} t_{\text{ind}}(F, G_n) = \mathbb{P}(\widehat{G}_n|_{[k]} = F) + o(1) \rightarrow \mathbb{P}(H|_{[k]} = F)$$

as $n \rightarrow \infty$, for every $F \in \mathcal{L}_k$, and Theorem 3.1 yields the existence of some random $\Gamma \in \mathcal{U}_\infty \subset \overline{\mathcal{U}}$ with $G_n \xrightarrow{d} \Gamma$ and $\mathbb{E} t_{\text{ind}}(F, \Gamma) = \mathbb{P}(H|_{[k]} = F)$. \square

5. EXCHANGEABLE RANDOM GRAPHS

Sexch

Definition. A random infinite graph $H \in \mathcal{L}_\infty$ is *exchangeable* if its distribution is invariant under every permutation of the vertices. (It is well-known that it is equivalent to consider only finite permutations, i.e., permutations σ of \mathbb{N} that satisfy $\sigma(i) = i$ for all sufficiently large i , so σ may be regarded as a permutation in \mathfrak{S}_n for some n .)

Equivalently, if $X_{ij} := \mathbf{1}[ij \in H]$ is the indicator of there being an edge ij in H , then the array $\{X_{ij}\}$, $1 \leq i, j \leq \infty$, is (jointly) exchangeable as defined in Section 1.

SJ

SJ

LE **Lemma 5.1.** *Let H be a random infinite graph in \mathcal{L}_∞ . Then the following are equivalent.*

- (i) H is exchangeable.
- (ii) $H|_{[k]}$ has a distribution invariant under all permutations of $[k]$, for every $k \geq 1$.
- (iii) $\mathbb{P}(H|_{[k]} = F)$ depends only on the isomorphism type of F , and can thus be seen as a function of F as an unlabelled graph in \mathcal{U}_k , for every $k \geq 1$.

Proof. (i) \implies (ii). Immediate.

(ii) \implies (i). If σ is a finite permutation of \mathbb{N} , then σ restricts to a permutation of $[k]$ for every large k , and it follows that if $H \circ \sigma$ is H with the vertices permuted by σ , then, for all large k $H \circ \sigma|_{[k]} = H|_{[k]} \circ \sigma \stackrel{d}{=} H|_{[k]}$, which implies $H \circ \sigma \stackrel{d}{=} H$.

(ii) \iff (iii). Trivial. □

TC1E **Theorem 5.2.** *The limit H is Theorem 4.1 is exchangeable.*

Proof. H satisfies Lemma 5.1(iii). □

Moreover, Theorem 4.1 implies the following connection with random elements of \mathcal{U}_∞ .

TE **Theorem 5.3.** *There is a one-to-one correspondence between distributions of random elements $\Gamma \in \mathcal{U}_\infty$ (or $\overline{\mathcal{U}}^*$) and distributions of exchangeable random infinite graphs $H \in \mathcal{L}_\infty$ given by*

$$\mathbb{E} t_{\text{ind}}(F, \Gamma) = \mathbb{P}(H|_{[k]} = F) \tag{5.1} \quad \text{e2a}$$

for every $k \geq 1$ and every $F \in \mathcal{L}_k$, or, equivalently,

$$\mathbb{E} t(F, \Gamma) = \mathbb{P}(H \supset F) \tag{5.2} \quad \text{e2b}$$

for every $F \in \mathcal{L}$. Furthermore, $H|_{[n]} \xrightarrow{d} \Gamma$ in $\overline{\mathcal{U}}$ as $n \rightarrow \infty$.

Proof. Note first that (5.1) and (5.2) are equivalent by (2.5) and (2.6), since $t(F, \Gamma) = t_{\text{inj}}(F, \Gamma)$ by (2.10), and $H \supset F$ if and only if $H|_{[k]} \supseteq F$ when $F \in \mathcal{L}_k$.

Suppose that Γ is a random element of $\mathcal{U}_\infty \subset \overline{\mathcal{U}}$. Since \mathcal{U} is dense in $\overline{\mathcal{U}}$, there exist (as in the proof of Corollary 3.3) random unlabelled graphs G_n such that $G_n \xrightarrow{\text{a.s.}} \Gamma$ in $\overline{\mathcal{U}}$ and thus $v(G_n) \xrightarrow{\text{a.s.}} \infty$ and $G_n \xrightarrow{d} \Gamma$. Hence, Theorems 4.1 and 5.2 show that $\widehat{G}_n \xrightarrow{d} H$ for some random exchangeable infinite graph H satisfying (5.1). Furthermore, (5.1) determines the distribution of $H|_{[k]}$ for every k , and thus the distribution of k .

Conversely, if H is an exchangeable random infinite graph, let $G_n = H|_{[n]}$. By Lemma 5.1(ii), the distribution of each G_n is invariant under permutations of the vertices, so if \widehat{G}_n is G_n with a random (re)labelling, we have $\widehat{G}_n \stackrel{d}{=} G_n$. Since $G_n \xrightarrow{d} H$ in \mathcal{L}_∞ (because \mathcal{L}_∞ has a product

topology), we thus have $\widehat{G}_n \xrightarrow{d} H$ in \mathcal{L}_∞ , so Theorem 4.1 applies and shows the existence of a random $\Gamma \in \mathcal{U}_\infty$ such that $G_n \xrightarrow{d} \Gamma$ and (5.1) holds. Finally (5.1) determines the distribution of Γ by Corollary 3.3. \square

Ras **Remark 5.1.** Moreover, $H|_{[n]}$ converges a.s. to some random variable $\Gamma \in \mathcal{U}_\infty$, because $t_{\text{ind}}(F, H|_{[n]})$, $n \geq v(F)$, is a reverse martingale for every $F \in \Gamma$. Alternatively, this follows by concentration estimates from the representation in Section 6, see Lovász and Szegedy [19, Theorem 2.5].

CE **Corollary 5.4.** *There is a one-to-one correspondence between elements Γ of $\mathcal{U}_\infty \cong \overline{\mathcal{U}}^*$ and extreme points of the set of distributions of exchangeable random infinite graphs $H \in \mathcal{L}_\infty$. This correspondence is given by*

$$t(F, \Gamma) = \mathbb{P}(H \supset F) \quad (5.3) \quad \text{ce}$$

for every $F \in \mathcal{L}$. Furthermore, $H|_{[n]} \xrightarrow{\text{a.s.}} \Gamma$ in $\overline{\mathcal{U}}$ as $n \rightarrow \infty$.

Proof. The extreme points of the set of distributions on \mathcal{U}_∞ are the point masses, which are in one-to-one correspondence with the elements of \mathcal{U}_∞ . \square

We can characterize these extreme point distributions of exchangeable random infinite graphs as follows.

TE2 **Theorem 5.5.** *Let H be an exchangeable random infinite graph. Then the following are equivalent.*

te2a (i) *The distribution of H is an extreme point in the set of exchangeable distributions in \mathcal{L}_∞ .*

te2b' (ii) *If F_1 and F_2 are two (finite) graphs with disjoint vertex sets $V(F_1), V(F_2) \subset \mathbb{N}$, then*

$$\mathbb{P}(H \supset F_1 \cup F_2) = \mathbb{P}(H \supset F_1) \mathbb{P}(H \supset F_2).$$

te2b (iii) *The restrictions $H|_{[k]}$ and $H|_{[k+1, \infty]}$ are independent for every k .*

te2c (iv) *Let \mathcal{F}_n be the σ -field generated by $H|_{[n, \infty]}$. Then the tail σ -field $\bigcap_{n=1}^\infty \mathcal{F}_n$ is trivial, i.e., contains only events with probability 0 or 1.*

Proof. (i) \implies (ii). By Corollary 5.4, H corresponds to some (non-random) $\Gamma \in \mathcal{U}_\infty$ such that

$$\mathbb{P}(H \supset F) = t(F, \Gamma) \quad (5.4) \quad \text{e6}$$

for every $F \in \mathcal{L}$. We have defined \mathcal{L} such that a graph $F \in \mathcal{L}$ is labelled by $1, \dots, v(F)$, but both sides of (5.4) are invariant under relabelling of F by arbitrary positive integers; the left hand side because H is exchangeable and the right hand side because $t(F, \Gamma)$ only depends on F as an unlabelled graph. Hence (5.4) holds for every finite graph F with $V(F) \subset \mathbb{N}$.

Furthermore, since Γ is non-random, Corollary 3.3 yields $t(F_1 \cup F_2, \Gamma) = t(F_1, \Gamma)t(F_2, \Gamma)$. Hence,

$$\mathbb{P}(H \supset F_1 \cup F_2) = t(F_1 \cup F_2, \Gamma) = t(F_1, \Gamma)t(F_2, \Gamma) = \mathbb{P}(H \supset F_1) \mathbb{P}(H \supset F_2).$$

(ii) \implies (iii). By inclusion–exclusion, as for (2.3), (ii) implies that if $1 \leq k < l < \infty$, then for any graphs F_1 and F_2 with $V(F_1) = \{1, \dots, k\}$

and $V(F_2) = \{k + 1, \dots, k + l\}$, the events $H|_{[k]} = F_1$ and $H|_{\{k+1, \dots, l\}} = F_2$ are independent. Hence $H|_{[k]}$ and $H|_{\{k, \dots, l\}}$ are independent for every $l > k$, and the result follows.

(iii) \implies (iv). Suppose A is an event in the tail σ -field $\bigcap_{n=1}^{\infty} \mathcal{F}_n$. Let \mathcal{F}_n^* be the σ -field generated by $H|_{[n]}$. By (iii), A is independent of \mathcal{F}_n^* for every n , and thus of the σ -field \mathcal{F} generated by $\bigcup \mathcal{F}_n^*$, which equals the σ -field \mathcal{F}_1 generated by H . However, $A \in \mathcal{F}_1$, so A is independent of itself and thus $\mathbb{P}(A) = 0$ or 1 .

(iv) \implies (i). Let $F \in \mathcal{L}_k$ for some k and let F_n be F with all vertices shifted by n . Consider the two indicators $I = \mathbf{1}[H \supseteq F]$ and $I_n = \mathbf{1}[H \supseteq F_n]$. Since I_n is \mathcal{F}_n -measurable,

$$\mathbb{P}(H \supseteq F \cup F_n) = \mathbb{E}(II_n) = \mathbb{E}(\mathbb{E}(I | \mathcal{F}_n)I_n). \tag{5.5} \quad \boxed{\text{e70}}$$

Moreover, $\mathbb{E}(I | \mathcal{F}_n)$, $n = 1, 2, \dots$, is a reverse martingale, and thus a.s. SJ

$$\mathbb{E}(I | \mathcal{F}_n) \rightarrow \mathbb{E}\left(I \mid \bigcap_{n=1}^{\infty} \mathcal{F}_n\right) = \mathbb{E}I,$$

using (iv). Hence, $(\mathbb{E}(I | \mathcal{F}_n) - \mathbb{E}I)I_n \rightarrow 0$ a.s., and by dominated convergence SJ

$$\mathbb{E}\left((\mathbb{E}(I | \mathcal{F}_n) - \mathbb{E}I)I_n\right) \rightarrow 0.$$

Consequently, (5.5) yields

$$\mathbb{P}(H \supseteq F \cup F_n) = \mathbb{E}I \mathbb{E}I_n + o(1) = \mathbb{P}(H \supseteq F) \mathbb{P}(H \supseteq F_n) + o(1).$$

Moreover, since H is exchangeable, $\mathbb{P}(H \supseteq F \cup F_n)$ (for $n \geq v(F)$) and $\mathbb{P}(H \supseteq F_n)$ do not depend on n , and we obtain as $n \rightarrow \infty$

$$\mathbb{P}(H \supseteq F \cup F_k) = \mathbb{P}(H \supseteq F)^2. \tag{5.6} \quad \boxed{\text{e7}}$$

Let Γ be a random element of \mathcal{U}_{∞} corresponding to H as in Theorem 5.3. By (5.2) and (3.2), (5.6) can be written

$$\mathbb{E}t(F, \Gamma)^2 = (\mathbb{E}t(F, \Gamma))^2.$$

Hence the random variable $t(F, \Gamma)$ has variance 0 so it is a.s. constant. Since this holds for every $F \in \mathcal{L}$, it follows that Γ is a.s. constant, i.e., we can take Γ non-random, and (i) follows by Corollary 5.4. □

Srep

6. REPRESENTATIONS OF GRAPH LIMITS AND EXCHANGEABLE GRAPHS

As said in the introduction, the exchangeable infinite random graphs were characterized by Aldous [1] and Hoover [14], see also Kallenberg [15], and the graph limits in $\mathcal{U}_{\infty} \cong \bar{\mathcal{U}}^*$ were characterized in a very similar way by Lovász and Szegedy [19]. We can now make the connection between these two characterizations explicit.

Let \mathcal{W} be the set of all measurable functions $W : [0, 1]^2 \rightarrow [0, 1]$ and let \mathcal{W}_s be the subset of symmetric functions. For every $W \in \mathcal{W}_s$, we define

an infinite random graph $G(\infty, W) \in \mathcal{L}_\infty$ as follows: we first choose a sequence X_1, X_2, \dots of i.i.d. random variables uniformly distributed on $[0, 1]$, and then, given this sequence, for each pair (i, j) with $i < j$ we draw an edge ij with probability $W(X_i, X_j)$, independently for all pairs (i, j) with $i < j$ (conditionally given $\{X_i\}$). Further, let $G(n, W)$ be the restriction $G(\infty, W)|_{[n]}$, which is obtained by the same construction with a finite sequence X_1, \dots, X_n .

It is evident that $G(\infty, W)$ is an exchangeable infinite random graph, and the result by Aldous and Hoover is that every exchangeable infinite random graph is obtained as a mixture of such $G(\infty, W)$; in other words as $G(\infty, W)$ with a random W .

Considering again a deterministic $W \in \mathcal{W}_s$, it is evident that Theorem 5.5(ii) holds, and thus Theorem 5.5 and Corollary 5.4 show that $G(\infty, W)$ corresponds to an element $\Gamma_W \in \mathcal{U}_\infty$. Moreover, by Theorem 5.3 and Remark 5.1, $G(n, W) \rightarrow \Gamma_W$ a.s. as $n \rightarrow \infty$, and (5.3) shows that if $F \in \mathcal{L}_k$, then

$$t(F, \Gamma_W) = \mathbb{P}(F \subseteq G(k, W)) = \int_{[0,1]^k} \prod_{ij \in \mathbb{E}(F)} W(x_i, x_j) dx_1 \dots dx_k. \quad (6.1) \quad \boxed{\text{tuw}}$$

The main result of Lovász and Szegedy [19] is that every element of $\mathcal{U}_\infty \cong \overline{\mathcal{U}^*}$ can be obtained as Γ_W satisfying (6.1) for some $W \in \mathcal{W}_s$.

It is now clear that the representation theorems of Aldous–Hoover [1, 14] and Lovász and Szegedy [19] are connected by Theorem 5.3 and Corollary 5.4 above, and that one characterization easily follows from the other.

Remark 6.1. The representations by W are far from unique, see Section 7. Borgs, Chayes, Lovász, Sós and Vesztegombi [8] call an element $W \in \mathcal{W}_s$ a *graphon*. They further define a pseudometric (called the *cut-distance*) on \mathcal{W}_s and show that if we consider the quotient space $\widehat{\mathcal{W}}_s$ obtained by identifying elements with cut-distance 0, we obtain a compact metric space, and the mapping $W \mapsto \Gamma_W$ yields a bijection $\widehat{\mathcal{W}}_s \rightarrow \overline{\mathcal{U}^*} \cong \mathcal{U}_\infty$, which furthermore is a homeomorphism.

Remark 6.2. As remarked in Lovász and Szegedy [19], we can more generally consider a symmetric measurable function $W : \mathcal{S}^2 \rightarrow [0, 1]$ for any probability space (\mathcal{S}, μ) , and define $G(\infty, W)$ as above with X_i i.i.d. random variables in \mathcal{S} with distribution μ . This does not give any new limit objects $G(\infty, W)$ or Γ_W , since we just said that every limit object is obtained from some $W \in \mathcal{W}_s$, but they can sometimes give useful representations.

An interesting case is when W is the adjacency matrix of a (finite) graph G , with $\mathcal{S} = V(G)$ and μ the uniform measure on \mathcal{S} ; we thus let X_i be i.i.d. random vertices of G and $G(n, W)$ equals the random graph $G[n]$ defined in Section 2. It follows from (6.1) and (2.1) that $t(F, \Gamma_W) = t(F, G)$ for every $F \in \mathcal{U}$, and thus $\Gamma_W = G$ as elements of $\overline{\mathcal{U}^*}$. In other words, $\Gamma_W \in \mathcal{U}_\infty = \pi(G)$, the “ghost” of G in $\mathcal{U}_\infty \cong \overline{\mathcal{U}^*}$.

Remark 6.3. For the asymptotic behavior of $G(n, W)$ in another, sparse, case, with W depending on n , see [3].

7. NON-UNIQUENESS

Sunique

SJ

The functions W on $[0, 1]^2$ used to represent graph limits or exchangeable arrays are far from unique. (For a special case when there is a natural canonical choice, which much simplifies and helps applications, see [12].) For example, it is obvious that if $\varphi : [0, 1] \rightarrow [0, 1]$ is any measure preserving map, then W and $W \circ \varphi$, defined by $W \circ \varphi(x, y) := W(\varphi(x), \varphi(y))$, define the same graph limit and the same (in distribution) exchangeable array.

Although in principle, this is the only source on non-uniqueness, the details are more complicated, mainly because the measure preserving map φ does not have to be a bijection, and thus the relation $W' = W \circ \varphi$ is not symmetric: it can hold without there being a measure preserving map φ' such that $W = W' \circ \varphi'$. (For a 1-dimensional example, consider $f(x) = x$ and $f'(x) = \varphi(x) = 2x \bmod 1$; for a 2-dimensional example, let $W(x, y) = f(x)f(y)$ and $W'(x, y) = f'(x)f'(y)$.)

For exchangeable arrays, the equivalence problem was solved by Hoover [14], who gave a criterion which in our case reduces to (vi) below; this criterion involves an auxiliary variable, and can be interpreted as saying $W = W' \circ \varphi'$ for a random φ' . This work was continued by Kallenberg, see [15], who gave a probabilistic proof and added criterion (v). For graph limits, Borgs, Chayes, Lovász, Sós and Vesztergombi [8] gave the criterion (vii) in terms of the cut-distance, and Bollobás and Riordan [4] found the criterion (v) in this context. Further, Borgs, Chayes, Lovász, Sós and Vesztergombi [8] announced the related criterion that there exists a measurable function $U : [0, 1]^2 \rightarrow [0, 1]$ and two measure preserving maps $\varphi, \varphi' : [0, 1] \rightarrow [0, 1]$ such that $W = U \circ \varphi$ and $W' = U \circ \varphi'$ a.e.; the proof of this will appear in [5].

As in Section 6, these two lines of work are connected by the results in Section 5, and we can combine the previous results as follows.

- Theorem 7.1.** *Let $W, W' \in \mathcal{W}_s$. Then the following are equivalent.*
- (i) $\Gamma_W = \Gamma_{W'}$ for the graph limits $\Gamma_W, \Gamma_{W'} \in \mathcal{U}_\infty$.
 - (ii) $t(F, \Gamma_W) = t(F, \Gamma_{W'})$ for every graph F .
 - (iii) The exchangeable random infinite graphs $G(\infty, W)$ and $G(\infty, W')$ have the same distribution.
 - (iv) The random graphs $G(n, W)$ and $G(n, W')$ have the same distribution for every finite n .
 - (v) There exist measure preserving maps $\varphi, \varphi' : [0, 1] \rightarrow [0, 1]$ such that $W \circ \varphi = W \circ \varphi'$ a.e. on $[0, 1]^2$, i.e., $W(\varphi(x), \varphi(y)) = W'(\varphi'(x), \varphi'(y))$ a.e.
 - (vi) There exists a measure preserving map $\psi : [0, 1]^2 \rightarrow [0, 1]$ such that $W(x_1, x_2) = W'(\psi(x_1, y_1), \psi(x_2, y_2))$ a.e. on $[0, 1]^4$.
 - (vii) $\delta_\square(W, W') = 0$, where δ_\square is the cut-distance defined in [8].

Proof. (i) \iff (ii). By our definition of $\mathcal{U}_\infty \subset \overline{\mathcal{U}}$.

(i) \iff (iii). By Corollary 5.4.

(iii) \iff (iv). Obvious.

(v) \implies (iii). If X_1, X_2, \dots are i.i.d. random variables uniformly distributed on $[0, 1]$, then so are $\varphi(X_1), \varphi(X_2), \dots$, and thus $G(\infty, W) \stackrel{d}{=} G(\infty, W \circ \varphi) = G(\infty, W' \circ \varphi') \stackrel{d}{=} G(\infty, W')$.

(iii) \implies (v). The general form of the representation theorem as stated in [15, Theorem 7.15, see also p. 304] is (in our two-dimensional case) $X_{ij} = f(\xi_\emptyset, \xi_i, \xi_j, \xi_{ij})$ for a function $f : [0, 1]^4 \rightarrow [0, 1]$, symmetric in the two middle variables, and independent random variables ξ_\emptyset, ξ_i ($1 \leq i$) and ξ_{ij} ($1 \leq i < j$), all uniformly distributed on $[0, 1]$, and where we further let $\xi_{ji} = \xi_{ij}$ for $j > i$. We can write the construction of $G(\infty, W)$ in this form with

$$f(\xi_\emptyset, \xi_i, \xi_j, \xi_{ij}) = \mathbf{1}[\xi_{ij} \leq W(\xi_i, \xi_j)]. \quad (7.1) \quad \boxed{\text{fW}}$$

Note that this f does not depend on ξ_\emptyset . (In general, ξ_\emptyset is needed for the case of a random W , which can be written as a deterministic function of ξ_\emptyset , but this is not needed in the present theorem.)

Suppose that $G(\infty, W) \stackrel{d}{=} G(\infty, W')$, let f be given by W by (7.1), and let similarly f' be given by W' ; for notational convenience we write $W_1 := W$, $W_2 := W'$, $f_1 := f$ and $f_2 := f'$. The equivalence theorem [15, Theorem 7.28] takes the form, using (7.1), that there exist measurable functions $g_{k,0} : [0, 1] \rightarrow [0, 1]$, $g_{k,1} : [0, 1]^2 \rightarrow [0, 1]$ and $g_{k,2} : [0, 1]^4 \rightarrow [0, 1]$, for $k = 1, 2$, that are measure preserving in the last coordinate for any fixed values of the other coordinates, and such that the two functions (for $k = 1$ and $k = 2$)

$$\begin{aligned} & f_k(g_{k,0}(\xi_\emptyset), g_{k,1}(\xi_\emptyset, \xi_1), g_{k,1}(\xi_\emptyset, \xi_2), g_{k,2}(\xi_\emptyset, \xi_1, \xi_2, \xi_{12})) \\ &= \mathbf{1}[W_k(g_{k,1}(\xi_\emptyset, \xi_1), g_{k,1}(\xi_\emptyset, \xi_2)) \geq g_{k,2}(\xi_\emptyset, \xi_1, \xi_2, \xi_{12})] \end{aligned}$$

are a.s. equal. Conditioned on ξ_\emptyset, ξ_1 and ξ_2 , the random variable $g_{k,2}(\xi_\emptyset, \xi_1, \xi_2, \xi_{12})$ is uniformly distributed on $[0, 1]$, and it follows (e.g., by taking the conditional expectation) that a.s.

$$W_1(g_{1,1}(\xi_\emptyset, \xi_1), g_{1,1}(\xi_\emptyset, \xi_2)) = W_2(g_{2,1}(\xi_\emptyset, \xi_1), g_{2,1}(\xi_\emptyset, \xi_2)).$$

For a.e. value x_0 of ξ_\emptyset , this thus holds for a.e. values of ξ_1 and ξ_2 , and we may choose $\varphi(x) = g_{1,1}(x_0, x)$ and $\varphi'(x) := g_{2,1}(x_0, x)$ for some such x_0 .

(iii) \iff (vi). Similar, using [15, Theorem 7.28(iii)].

(ii) \iff (vii). See [8]. \square

Sbip

8. BIPARTITE GRAPHS

The definitions and results above have analogues for bipartite graphs, which we give in this section, leaving some details to the reader. The proofs are straightforward analogues of the ones given above and are omitted. Applications of the results of this section to random difference graphs are in [12].

A *bipartite graph* will be a graph with an explicit bipartition; in other words, a bipartite graph G consists of two vertex sets $V_1(G)$ and $V_2(G)$ and an edge set $E(G) \subseteq V_1(G) \times V_2(G)$; we let $v_1(G) := |V_1(G)|$ and $v_2(G) := |V_2(G)|$ be the numbers of vertices in the two sets. Again we consider both the labelled and unlabelled cases; in the labelled case we assume the labels of the vertices in $V_j(G)$ are $1, \dots, v_j(G)$ for $j = 1, 2$. Let $\mathcal{B}_{n_1 n_2}^L$ be the set of the $2^{n_1 n_2}$ labelled bipartite graphs with vertex sets $[n_1]$ and $[n_2]$, and let $\mathcal{B}_{n_1 n_2}$ be the quotient set $\mathcal{B}_{n_1 n_2}^L / \cong$ of unlabelled bipartite graphs with n_1 and n_2 vertices in the two parts; further, let $\mathcal{B}^L := \bigcup_{n_1, n_2 \geq 1} \mathcal{B}_{n_1 n_2}^L$ and $\mathcal{B} := \bigcup_{n_1, n_2 \geq 1} \mathcal{B}_{n_1 n_2}$.

We let $G[k_1, k_2]$ be the random graph in $\mathcal{B}_{k_1 k_2}^L$ obtained by sampling k_j vertices from $V_j(G)$ ($j = 1, 2$), uniformly with replacement, and let, provided $k_j \leq v_j(G)$, $G[k_1, k_2]'$ be the corresponding random graph obtained by sampling without replacement. We then define $t(F, G)$, $t_{\text{inj}}(F, G)$ and $t_{\text{ind}}(F, G)$ for (unlabelled) bipartite graphs F and G in analogy with (2.1)–(2.3). Then (2.4)–(2.6) still hold, *mutatis mutandis*; for example,

$$|t(F, G) - t_{\text{inj}}(F, G)| \leq \frac{v_1(F)^2}{2v_1(G)} + \frac{v_2(F)^2}{2v_2(G)}. \quad (8.1) \quad \boxed{\text{a4xx}}$$

In analogy with (2.7), we now define $\tau : \mathcal{B} \rightarrow [0, 1]^{\mathcal{B}}$ by

$$\tau(G) := (t(F, G))_{F \in \mathcal{B}} \in [0, 1]^{\mathcal{B}}. \quad (8.2) \quad \boxed{\text{taub}}$$

We define $\mathcal{B}^* := \tau(\mathcal{B}) \subseteq [0, 1]^{\mathcal{B}}$ to be the image of \mathcal{B} under this mapping τ , and let $\overline{\mathcal{B}^*}$ be the closure of \mathcal{B}^* in $[0, 1]^{\mathcal{B}}$; this is a compact metric space.

Again, τ is not injective; we may consider a graph G as an element of \mathcal{B}^* by identifying G and $\tau(G)$, but this implies identification of some graphs of different orders and we prefer to avoid it. We let \mathcal{B}^+ be the union of \mathcal{B} and some two-point set $\{*_1, *_2\}$ and consider the mapping $\tau^+ : \mathcal{B} \rightarrow [0, 1]^{\mathcal{B}^+} = [0, 1]^{\mathcal{B}} \times [0, 1] \times [0, 1]$ defined by

$$\tau^+(G) = (\tau(G), v_1(G)^{-1}, v_2(G)^{-1}). \quad (8.3) \quad \boxed{\text{tab+}}$$

Then τ^+ is injective and we can identify \mathcal{B} with its image $\tau^+(\mathcal{B}) \subseteq [0, 1]^{\mathcal{B}^+}$ and define $\overline{\mathcal{B}} \subseteq [0, 1]^{\mathcal{B}^+}$ as its closure; this is a compact metric space.

The functions $t(F, \cdot)$, $t_{\text{inj}}(F, \cdot)$, $t_{\text{ind}}(F, \cdot)$ and $v_j(\cdot)^{-1}$, for $F \in \mathcal{B}$ and $j = 1, 2$, have unique continuous extensions to $\overline{\mathcal{B}}$.

We let $\mathcal{B}_{\infty\infty} := \{G \in \overline{\mathcal{B}} : v_1(G) = v_2(G) = \infty\}$; this is the set of all limit objects of sequences (G_n) in \mathcal{B} with $v_1(G_n), v_2(G_n) \rightarrow \infty$. By (8.1), $t_{\text{inj}}(F, G) = t(F, G)$ for every $G \in \mathcal{B}_{\infty\infty}$ and every $F \in \mathcal{B}$. The projection $\pi : \overline{\mathcal{B}} \rightarrow \overline{\mathcal{B}^*}$ restricts to a homeomorphism $\mathcal{B}_{\infty\infty} \cong \overline{\mathcal{B}^*}$.

Remark 8.1. Note that in the bipartite case there are other limit objects too in $\overline{\mathcal{B}}$; in fact, $\overline{\mathcal{B}}$ can be partitioned into \mathcal{B} , $\mathcal{B}_{\infty\infty}$, and the sets $\mathcal{B}_{n\infty}$, $\mathcal{B}_{\infty n}$, for $n = 1, 2, \dots$, where, for example, $\mathcal{B}_{n_1\infty}$ is the set of limits of sequences (G_n) of bipartite graphs such that $v_2(G_n) \rightarrow \infty$ but $v_1(G_n) = n_1$ is constant. We will not consider such degenerate limits further here, but we remark that

in the simplest case $n_1 = 1$, a bipartite graph in $\mathcal{B}_{1n_2}^L$ can be identified with a subset of $[n_2]$, and an unlabelled graph in \mathcal{B}_{1n_2} thus with a number in $m \in \{0, \dots, n_2\}$, the number of edges in the graph, and it is easily seen that a sequence of such unlabelled graphs with $n_2 \rightarrow \infty$ converges in $\overline{\mathcal{B}}$ if and only if the proportion m/n_2 converges; hence we can identify $\mathcal{B}_{1\infty}$ with the interval $[0,1]$.

We have the following basic result, cf. Theorem 2.1.

T1B **Theorem 8.1.** *Let (G_n) be a sequence of bipartite graphs with $v_1(G_n), v_2(G_n) \rightarrow \infty$. Then the following are equivalent.*

- T1Bt** (i) $t(F, G_n)$ converges for every $F \in \mathcal{B}$.
- T1Btinj** (ii) $t_{\text{inj}}(F, G_n)$ converges for every $F \in \mathcal{B}$.
- T1Btind** (iii) $t_{\text{ind}}(F, G_n)$ converges for every $F \in \mathcal{B}$.
- (iv) G_n converges in $\overline{\mathcal{B}}$.

In this case, the limit G of G_n belongs to $\mathcal{B}_{\infty\infty}$ and the limits in (i), (iii) and (iv) are $t(F, G)$, $t_{\text{inj}}(F, G)$ and $t_{\text{ind}}(F, G)$.

For convergence of random unlabelled bipartite graphs, the results in Section 3 hold with trivial changes.

T2B **Theorem 8.2.** *Let $G_n, n \geq 1$, be random unlabelled bipartite graphs and assume that $v_1(G_n), v_2(G_n) \xrightarrow{P} \infty$. The following are equivalent, as $n \rightarrow \infty$.*

- T2Ba** (i) $G_n \xrightarrow{d} \Gamma$ for some random $\Gamma \in \overline{\mathcal{B}}$.
- T2Bb** (ii) *For every finite family F_1, \dots, F_m of (non-random) bipartite graphs, the random variables $t(F_1, G_n), \dots, t(F_m, G_n)$ converge jointly in distribution.*
- T2Bc** (iii) *For every (non-random) $F \in \mathcal{B}$, the random variables $t(F, G_n)$ converge in distribution.*
- T2Bd** (iv) *For every (non-random) $F \in \mathcal{B}$, the expectations $\mathbb{E}t(F, G_n)$ converge.*

If these properties hold, then the limits in (ii), (iii) and (iv) are $(t(F_i, \Gamma))_{i=1}^m, t(F, \Gamma)$ and $\mathbb{E}t(F, \Gamma)$, respectively. Furthermore, $\Gamma \in \mathcal{B}_{\infty\infty}$ a.s.

The same results hold if t is replaced by t_{inj} or t_{ind} .

C2B **Corollary 8.3.** *Let $G_n, n \geq 1$, be random unlabelled bipartite graphs such that $v_1(G_n), v_2(G_n) \xrightarrow{P} \infty$, and let $G \in \mathcal{B}_{\infty\infty}$. The following are equivalent, as $n \rightarrow \infty$.*

- C2Ba** (i) $G_n \xrightarrow{P} G$.
- C2Bc** (ii) $t(F, G_n) \xrightarrow{P} t(F, G)$ for every (non-random) $F \in \mathcal{B}$.
- C2Bd** (iii) $\mathbb{E}t(F, G_n) \rightarrow t(F, G)$ for every (non-random) $F \in \mathcal{B}$.

The same result holds if t is replaced by t_{inj} or t_{ind} .

As above, the distribution of Γ is uniquely determined by the numbers $\mathbb{E}t(F, \Gamma), F \in \mathcal{B}$.

Let $\mathcal{B}_{\infty\infty}^L$ denote the set of all labelled infinite bipartite graphs with the vertex sets $V_1(G) = V_2(G) = \mathbb{N}$. $\mathcal{B}_{\infty\infty}^L$ is a compact metric space with the natural product topology.

If G is a bipartite graph, let \widehat{G} be the random labelled bipartite graph obtained by random labellings of the vertices in $V_j(G)$ by the numbers $1, \dots, v_j(G)$, for $j = 1, 2$. This is a random finite bipartite graph, but we can also regard it as a random element of $\mathcal{B}_{\infty\infty}^L$ by adding isolated vertices.

Definition. A random infinite bipartite graph $H \in \mathcal{B}_{\infty\infty}^L$ is *exchangeable* if its distribution is invariant under every pair of finite permutations of $V_1(H)$ and $V_2(H)$.

TC1B **Theorem 8.4.** *Let (G_n) be a sequence of random graphs in \mathcal{B} and assume that $v_1(G_n), v_2(G_n) \xrightarrow{p} \infty$. Then the following are equivalent.*

- (i) $G_n \xrightarrow{d} \Gamma$ in $\overline{\mathcal{B}}$ for some random $\Gamma \in \overline{\mathcal{B}}$.
- (ii) $\widehat{G_n} \xrightarrow{d} H$ in $\mathcal{B}_{\infty\infty}^L$ for some random $H \in \mathcal{B}_{\infty\infty}^L$.

If these hold, then $\mathbb{P}(H|_{[k_1] \times [k_2]} = F) = \mathbb{E} t_{\text{ind}}(F, \Gamma)$ for every $F \in \mathcal{B}_{k_1 k_2}^L$. Furthermore, $\Gamma \in \mathcal{B}_{\infty\infty}$ a.s., and H is exchangeable.

TEB **Theorem 8.5.** *There is a one-to-one correspondence between distributions of random elements $\Gamma \in \mathcal{B}_{\infty\infty}$ (or $\overline{\mathcal{B}^*}$) and distributions of exchangeable random infinite graphs $H \in \mathcal{B}_{\infty\infty}^L$ given by*

$$\mathbb{E} t_{\text{ind}}(F, \Gamma) = \mathbb{P}(H|_{[k_1] \times [k_2]} = F) \quad (8.4)$$

for every $k_1, k_2 \geq 1$ and every $F \in \mathcal{B}_{k_1 k_2}^L$, or, equivalently,

$$\mathbb{E} t(F, \Gamma) = \mathbb{P}(H \supset F) \quad (8.5)$$

for every $F \in \mathcal{B}^L$. Furthermore, $H|_{[n_1] \times [n_2]} \xrightarrow{d} \Gamma$ in $\overline{\mathcal{B}}$ as $n_1, n_2 \rightarrow \infty$.

CEB **Corollary 8.6.** *There is a one-to-one correspondence between elements Γ of $\mathcal{B}_{\infty\infty} \cong \overline{\mathcal{B}^*}$ and extreme points of the set of distributions of exchangeable random infinite graphs $H \in \mathcal{B}_{\infty\infty}^L$. This correspondence is given by*

$$t(F, \Gamma) = \mathbb{P}(H \supset F) \quad (8.6)$$

for every $F \in \mathcal{B}^L$. Furthermore, $H|_{[n_1] \times [n_2]} \xrightarrow{p} \Gamma$ in $\overline{\mathcal{B}}$ as $n_1, n_2 \rightarrow \infty$.

Remark 8.2. We have not checked whether $H|_{[n_1] \times [n_2]} \xrightarrow{\text{a.s.}} \Gamma$ in $\overline{\mathcal{B}}$ as $n_1, n_2 \rightarrow \infty$. This holds at least for a subsequence $(n_1(m), n_2(m))$ with both $n_1(m)$ and $n_2(m)$ non-decreasing because then $t_{\text{inj}}(F, H|_{[n_1] \times [n_2]})$ is a reverse martingale.

TE2B **Theorem 8.7.** *Let H be an exchangeable random infinite bipartite graph. Then the following are equivalent.*

- te2Ba** (i) *The distribution of H is an extreme point in the set of exchangeable distributions in $\mathcal{B}_{\infty\infty}^L$.*

te2Bb'

- (ii) If F_1 and F_2 are two (finite) bipartite graphs with the vertex sets $V_j(F_1)$ and $V_j(F_2)$ disjoint subsets of \mathbb{N} for $j = 1, 2$, then

$$\mathbb{P}(H \supset F_1 \cup F_2) = \mathbb{P}(H \supset F_1) \mathbb{P}(H \supset F_2).$$

The construction in Section 6 takes the following form; note that there is no need to assume symmetry of W . For every $W \in \mathcal{W}$, we define an infinite random bipartite graph $G(\infty, \infty, W) \in \mathcal{B}_{\infty\infty}^L$ as follows: we first choose two sequence X_1, X_2, \dots and Y_1, Y_2, \dots of i.i.d. random variables uniformly distributed on $[0, 1]$, and then, given these sequences, for each pair $(i, j) \in \mathbb{N} \times \mathbb{N}$ we draw an edge ij with probability $W(X_i, Y_j)$, independently for all pairs (i, j) . Further, let $G(n_1, n_2, W)$ be the restriction $G(\infty, \infty, W)|_{[n_1] \times [n_2]}$, which is obtained by the same construction with finite sequences X_1, \dots, X_{n_1} and Y_1, \dots, Y_{n_2} .

It is evident that $G(\infty, \infty, W)$ is an exchangeable infinite random bipartite graph. Furthermore, it satisfies Theorem 8.7(ii). Theorem 8.5 and Corollary 8.6 yield a corresponding element $\Gamma_W'' \in \mathcal{B}_{\infty\infty} \cong \overline{\mathcal{B}}^*$ such that $G(n_1, n_2, W) \xrightarrow{P} \Gamma_W''$ as $n_1, n_2 \rightarrow \infty$ and, for every $F \in \mathcal{B}_{k_1 k_2}^L$,

$$t(F, \Gamma_W'') = \int_{[0,1]^{k_1+k_2}} \prod_{ij \in \mathbb{E}(F)} W(x_i, y_j) dx_1 \dots dx_{k_1} dy_1 \dots dy_{k_2}. \quad (8.7) \quad \text{tuww}$$

The result by Aldous [1] in the non-symmetric case is that every exchangeable infinite random bipartite graph is obtained as a mixture of such $G(\infty, \infty, W)$; in other words as $G(\infty, \infty, W)$ with a random W .

By Theorem 8.5 and Corollary 8.6 above, this implies (and is implied by) the fact that every element of $\overline{\mathcal{B}}$ equals Γ_W'' for some (non-unique) $W \in \mathcal{W}$; the bipartite version of the characterization by Lovász and Szegedy [19].

9. DIRECTED GRAPHS

SJ

Sdir

A *directed graph* G consists of a vertex set $V(G)$ and an edge set $E(G) \subseteq V(G) \times V(G)$; the edge indicators thus form an arbitrary zero–one matrix $\{X_{ij}\}$, $i, j \in V(G)$. Note that we allow loops, corresponding to the diagonal indicators X_{ii} . The definitions and results above have analogues for directed graphs too, with mainly notational differences. We sketch these in this section, leaving the details to the reader.

Let \mathcal{D}_n^L be the set of the 2^{n^2} labelled directed graphs with vertex set $[n]$ and let \mathcal{D}_n be the quotient set \mathcal{D}_n^L / \cong of unlabelled directed graphs with n vertices; further, let $\mathcal{D}^L := \bigcup_{n \geq 1} \mathcal{D}_n^L$ and $\mathcal{D} := \bigcup_{n \geq 1} \mathcal{D}_n$.

The definitions in Section 2 apply to directed graphs too, with at most notational differences. $G[k]$ and $G[k]'$ now are random directed graphs and $t(F, G)$, $t_{\text{inj}}(F, G)$ and $t_{\text{ind}}(F, G)$ are defined for (unlabelled) directed graphs F and G by (2.1)–(2.3). We now define $\tau : \mathcal{D} \rightarrow [0, 1]^{\mathcal{D}}$ by, cf. (2.7),

$$\tau(G) := (t(F, G))_{F \in \mathcal{D}} \in [0, 1]^{\mathcal{D}}. \quad (9.1) \quad \text{taud}$$

We define $\mathcal{D}^* := \tau(\mathcal{D}) \subseteq [0, 1]^{\mathcal{D}}$ to be the image of \mathcal{D} under this mapping τ , and let $\overline{\mathcal{D}^*}$ be the closure of \mathcal{D}^* in $[0, 1]^{\mathcal{D}}$; this is a compact metric space.

Again, τ is not injective. We let \mathcal{D}^+ be the union of \mathcal{D} and some one-point set $\{*\}$ and consider the mapping $\tau^+ : \mathcal{D} \rightarrow [0, 1]^{\mathcal{D}^+} = [0, 1]^{\mathcal{D}} \times [0, 1]$ defined by (2.9) as before. Then τ^+ is injective and we can identify \mathcal{D} with its image $\tau^+(\mathcal{D}) \subseteq [0, 1]^{\mathcal{D}^+}$ and define $\overline{\mathcal{D}} \subseteq [0, 1]^{\mathcal{D}^+}$ as its closure; this is a compact metric space. The functions $t(F, \cdot)$, $t_{\text{inj}}(F, \cdot)$, $t_{\text{ind}}(F, \cdot)$ and $v(\cdot)^{-1}$, for $F \in \mathcal{D}$, have unique continuous extensions to $\overline{\mathcal{D}}$.

We let $\mathcal{D}_\infty := \{G \in \overline{\mathcal{D}} : v(G) = \infty\}$; this is the set of all limit objects of sequences (G_n) in \mathcal{D} with $v(G_n) \rightarrow \infty$. By (8.1), $t_{\text{inj}}(F, G) = t(F, G)$ for every $G \in \mathcal{D}_\infty$ and every $F \in \mathcal{D}$. The projection $\pi : \overline{\mathcal{D}} \rightarrow \overline{\mathcal{D}^*}$ restricts to a homeomorphism $\mathcal{D}_\infty \cong \overline{\mathcal{D}^*}$.

All results in Sections 2–5 are valid for directed graphs too, with at most notational differences.

Please check!

The main difference for the directed case concerns the representations discussed in Section 6. Since two vertices may be connected by up to two directed edges (in opposite directions), and the events that the two possible edges occur typically are dependent, a single function W is no longer enough. Instead, we have a representation using several functions as follows.

Please check!!

Let \mathcal{W}_5 be the set of quintuples $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$ where $W_{\alpha\beta} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ are measurable functions such that $\sum_{\alpha, \beta=0}^1 W_{\alpha\beta} = 1$ and $W_{\alpha\beta}(x, y) = W_{\beta\alpha}(y, x)$ for $\alpha, \beta \in \{0, 1\}$ and $x, y \in [0, 1]$. For $\mathbf{W} \in \mathcal{W}_5$, we define a random infinite directed graph $G(\infty, \mathbf{W})$ by specifying its edge indicators X_{ij} as follows: we first choose a sequence Y_1, Y_2, \dots of i.i.d. random variables uniformly distributed on $[0, 1]$, and then, given this sequence, let $X_{ii} = w(Y_i)$ and for each pair (i, j) with $i < j$ choose X_{ij} and X_{ji} at random such that

$$\mathbb{P}(X_{ij} = \alpha \text{ and } X_{ji} = \beta) = W_{\alpha\beta}(Y_i, Y_j), \quad \alpha, \beta \in \{0, 1\}; \quad (9.2) \quad \boxed{\text{dirw}}$$

this is done independently for all pairs (i, j) with $i < j$ (conditionally given $\{Y_i\}$). In other words, for every i we draw a loop at i if $w(Y_i) = 1$ and for each pair (i, j) with $i < j$ we draw edges ij and ji at random such that (9.2) holds. Further, let $G(n, \mathbf{W})$ be the restriction $G(\infty, \mathbf{W})|_{[n]}$, which is obtained by the same construction with a finite sequence Y_1, \dots, Y_n .

In particular, note that the loops appear independently, each with probability $p = \mathbb{P}(w(Y_1) = 1)$. We may specify the loops more clearly by the following alternative version of the construction. Let $\mathcal{S} := [0, 1] \times \{0, 1\}$ and let \mathcal{W}_4 be the set of quadruples $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11})$ where $W_{\alpha\beta} : \mathcal{S}^2 \rightarrow [0, 1]$ are measurable functions such that $\sum_{\alpha, \beta=0}^1 W_{\alpha\beta} = 1$ and $W_{\alpha\beta}(x, y) = W_{\beta\alpha}(y, x)$ for $\alpha, \beta \in \{0, 1\}$ and $x, y \in \mathcal{S}$. For every $\mathbf{W} \in \mathcal{W}_4$ and $p \in [0, 1]$, we define a random infinite directed graph $G(\infty, \mathbf{W}, p)$ by specifying its edge indicators X_{ij} as follows: We first choose sequences ξ_1, ξ_2, \dots and ζ_1, ζ_2, \dots of random variables, all independent, with $\xi_i \sim U(0, 1)$ and $\zeta_i \sim \text{Be}(p)$, i.e., $\zeta_i \in \{0, 1\}$ with $\mathbb{P}(\zeta_i = 1) = p$; we let

$Y_i := (\xi_i, \zeta_i) \in \mathcal{S}$. Then, given these sequences, let $X_{ii} = \zeta_i$ and for each pair (i, j) with $i < j$ choose X_{ij} and X_{ji} at random according to (9.2), independently for all pairs (i, j) with $i < j$ (conditionally given $\{Y_i\}$). In other words, ζ_i is the indicator of a loop at i . Further, let $G(n, \mathbf{W}, p)$ be the restriction $G(\infty, \mathbf{W}, p)|_{[n]}$, which is obtained by the same construction with a finite sequence Y_1, \dots, Y_n .

It is obvious from the symmetry of the construction that the random infinite directed graphs $G(\infty, \mathbf{W})$ and $G(\infty, \mathbf{W}, p)$ are exchangeable. Further, using Theorem 5.5, their distributions are extreme points, so by Corollary 5.4 they correspond to directed graph limits, i.e., elements of \mathcal{D}_∞ , which we denote by $\Gamma_{\mathbf{W}}$ and $\Gamma_{\mathbf{W}, p}$, respectively; (5.3) shows that if $F \in \mathcal{D}_k$, then

$$t(F, \Gamma_{\mathbf{W}}) = \mathbb{P}(F \subseteq G(k, \mathbf{W})), \quad t(F, \Gamma_{\mathbf{W}, p}) = \mathbb{P}(F \subseteq G(k, \mathbf{W}, p)).$$

By Theorem 5.3 and Remark 5.1, $G(n, \mathbf{W}) \rightarrow \Gamma_{\mathbf{W}}$ and $G(n, \mathbf{W}, p) \rightarrow \Gamma_{\mathbf{W}, p}$ a.s. as $n \rightarrow \infty$.

We can show a version of the representation results in Section 6 for directed graphs.

TD **Theorem 9.1.** *An exchangeable random infinite directed graph is obtained as a mixture of $G(\infty, \mathbf{W})$; in other words, as $G(\infty, \mathbf{W})$ with a random \mathbf{W} . Alternatively, it is obtained as a mixture of $G(\infty, \mathbf{W}, p)$; in other words, as $G(\infty, \mathbf{W}, p)$ with a random (\mathbf{W}, p) .*

Every directed graph limit, i.e., every element of \mathcal{D}_∞ , is $\Gamma_{\mathbf{W}}$ for some $\mathbf{W} \in \mathcal{W}_5$, or equivalently $\Gamma_{\mathbf{W}, p}$ for some $\mathbf{W} \in \mathcal{W}_4$ and $p \in [0, 1]$.

Proof. For jointly exchangeable random arrays $\{X_{ij}\}$ of zero–one variables, the Aldous–Hoover representation theorem takes the form [15, Theorem 7.22]

$$\begin{aligned} X_{ii} &= f_1(\xi_\emptyset, \xi_i), \\ X_{ij} &= f_2(\xi_\emptyset, \xi_i, \xi_j, \xi_{ij}), \quad i \neq j, \end{aligned}$$

where $f_1 : [0, 1]^2 \rightarrow \{0, 1\}$ and $f_2 : [0, 1]^4 \rightarrow \{0, 1\}$ are two measurable functions, $\xi_{ji} = \xi_{ij}$, and ξ_\emptyset, ξ_i ($1 \leq i$) and ξ_{ij} ($1 \leq i < j$) are independent random variables uniformly distributed on $[0, 1]$ (as in the proof of Theorem 7.1). If further the distribution of the array $\{X_{ij}\}$ is an extreme point in the set of exchangeable distributions, then by Theorem 5.5 and [15, Lemma 7.35], there exists such a representation where f_1 and f_2 do not depend on ξ_\emptyset , so $X_{ii} = f_1(\xi_i)$ and $X_{ij} = f_2(\xi_i, \xi_j, \xi_{ij})$, $i \neq j$. In this case, define $w = f_1$ and

$$W_{\alpha\beta}(x, y) := \mathbb{P}(f_2(x, y, \xi) = \alpha \text{ and } f_2(y, x, \xi) = \beta), \quad \alpha, \beta \in \{0, 1\},$$

where $\xi \sim U(0, 1)$. This defines a quintuple $\mathbf{W} \in \mathcal{W}_5$, such that the edge indicators X_{ij} of $G(\infty, \mathbf{W})$ have the desired distribution.

In general, the variable ξ_\emptyset can be interpreted as making \mathbf{W} random.

To obtain the alternative representation, let $\zeta_i := w(\xi_i) = X_{ii}$ and $p := \mathbb{P}(\zeta_i = 1)$. There exists a measure preserving map $\phi : (\mathcal{S}, \mu_p) \rightarrow [0, 1]$ such

that $[0, 1] \times \{j\}$ is mapped onto $\{x \in [0, 1] : w(x) = j\}$ for $j = 0, 1$ (i.e., $w \circ \phi(x, \zeta) = \zeta$), and we can use the quadruple $(W_{\alpha\beta} \circ \phi)_{\alpha,\beta}$.

The representations for graph limits follow by Corollary 5.4 as discussed above. \square

Example 9.2. A *random tournament* T_n is a random directed graph on n vertices without loops where each pair of vertices is connected by exactly one edge, with random direction (with equal probabilities for the two directions, and independent of all other edges). This equals $G(n, \mathbf{W})$ or $G(n, \mathbf{W}, p)$ with $W_{00} = W_{11} = 0$, $W_{01} = W_{10} = 1/2$, and $w = 0$ or $p = 0$, and converges thus a.s. to the limit $\Gamma_{\mathbf{W},0}$ for $\mathbf{W} = (W_{\alpha\beta})_{\alpha,\beta}$.

Note that if $\{X_{ij}\}$ are the edge indicators of an exchangeable random infinite directed graph, then the loop indicators $\{X_{ii}\}$ form a binary exchangeable sequence, and the representation as $G(\infty, \mathbf{W}, p)$ in Theorem 9.1 exhibits them as a mixture of i.i.d. $\text{Be}(p)$ variable, which has brought us back to deFinetti's theorem 1.1.

REFERENCES

Aldous	[1] D. Aldous, Representations for partially exchangeable arrays of random variables. <i>J. Multivar. Anal.</i> 11 , 581–598, 1981.
Bill1	[2] P. Billingsley, <i>Convergence of Probability Measures</i> . Wiley, New York, 1968.
SJ178	[3] B. Bollobás, S. Janson and O. Riordan, The phase transition in inhomogeneous random graphs. <i>Random Struct. Alg.</i> 31 (2007), 3–122.
BR	[4] B. Bollobás and O. Riordan, Sparse graphs: metrics and random models. Preprint, 2007. http://arxiv.org/0708.1919v1
BCL:unique	[5] C. Borgs, J. T. Chayes and L. Lovász, Unique limits of dense graph sequences. In preparation.
BCL3	[6] C. Borgs, J. T. Chayes, L. Lovász, V. T. Sós, B. Szegedy and K. Vesztergombi, Graph limits and parameter testing. In <i>STOC'06: Proceedings of the 38th Annual ACM Symposium on Theory of Computing</i> , 261–270. ACM, New York, 2006.
BCL2	[7] C. Borgs, J. T. Chayes, L. Lovász, V. T. Sós and K. Vesztergombi, Counting graph homomorphisms. In <i>Topics in Discrete Mathematics</i> (ed. M. Klazar, J. Kratochvil, M. Loeb, J. Matousek, R. Thomas, P. Valtr), 315–371. Springer, New York, 2006.
BCL1	[8] C. Borgs, J. T. Chayes, L. Lovász, V. T. Sós and K. Vesztergombi, Convergent sequences of dense graphs I: Subgraph frequencies, metric properties and testing. Preprint, 2007. http://arxiv.org/math.CO/0702004
BCL1ii	[9] C. Borgs, J. T. Chayes, L. Lovász, V. T. Sós and K. Vesztergombi, Convergent sequences of dense graphs II: Multiway cuts and statistical physics. Preprint, 2007. http://research.microsoft.com/~borgs/
DF1981	[10] P. Diaconis and D. Freedman, On the statistics of vision: The Julesz conjecture. <i>J. Math. Psychol.</i> 24 , 112–138, 1981.

- DF1984** [11] P. Diaconis and D. Freedman, Partial exchangeability and sufficiency. In *Statistics: Applications and New Directions* (eds. J. K. Jhosh and J. Roy), 205–236. Indian Statistical Institute, Calcutta, 1984.
- threshold** [12] P. Diaconis, S. Holmes and S. Janson, Threshold graph limits and random threshold graphs. In preparation.
- Freed** [13] M. Freedman, L. Lovász and A. Schrijver, Reflection positivity, rank connectivity, and homomorphism of graphs. *J. Amer. Math. Soc.* **20**, 37–51, 2007.
- Hoover** [14] D. Hoover, *Relations on Probability Spaces and Arrays of Random Variables*. Preprint, Institute for Advanced Study, Princeton, NJ, 1979.
- Kallenberg:exch** [15] O. Kallenberg, *Probabilistic Symmetries and Invariance Principles*. Springer, New York, 2005.
- LL2006** [16] L. Lovász, The rank of connection matrices and the dimension of graph algebras. *Eur. J. Comb.* **27**, 962–970, 2006.
- LL2007** [17] L. Lovász, Connection matrices. In *Combinatorics, Complexity, and Chance: A Tribute to Dominic Welsh* (ed. G. Grimmet and C. McDiarmid), 179–190. Oxford University Press, Oxford, 2007.
- LSos** [18] L. Lovász and V. T. Sós, Generalized quasirandom graphs. *J. Comb. Theory B.* **98** (2008), Issue 1, 146–163.
- LSz** [19] L. Lovász and B. Szegedy, Limits of dense graph sequences. *J. Comb. Theory B* **96**, 933–957, 2006.
- LSzszem** [20] L. Lovász and B. Szegedy, Szemerédi’s Lemma for the analyst. *J. Geom. Func. Anal.* **17**, 252–270, 2007.
- LSzcont** [21] L. Lovász and B. Szegedy, Contractors and connectors in graph algebras. *J. Comb. Theory B*, to appear.
- LSztest** [22] L. Lovász and B. Szegedy, Testing properties of graphs and functions. Preprint, 2007.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD CALIFORNIA 94305, USA AND DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE NICE - SOPHIA ANTIPOLIS, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE

DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY, PO Box 480, SE-751 06 UPPSALA, SWEDEN

E-mail address: svante.janson@math.uu.se

URL: <http://www.math.uu.se/~svante/>